THE INNERVATION OF THE BAT COCHLEA

by

WILHELM FIRBAS

Anatomisches Institut, Währingerstr. 13, A 1090, Wien, Oesterreich

For different species of bats, fixed in 5 % formaldehyd, an estimation of the number of neurons in the spiral ganglion was made. The cochleae were decalcified in EDTA and embedded in paraffin. The complete series of sections were stained with hematoxylin. On the sections through the ganglion, the number of ganglion cell nuclei was determined, with the help of an eye-piece micrometer. A 200 times magnification was used. The figures obtained with this method were corrected, with the help

of Floderus' formula:
$$N = n \cdot \frac{T}{T + d - 2k} \quad (N =$$

the corrected estimate of the number of nuclei centres within the section; n = the observed count; T = section thickness; d = diameter of nuclei; k = vertical thickness of the smallest visible fragment). This formula helps to avoid double counting of split nuclei. After this correction the following results were obtained (table I).

Table 1. Number of nerve cells in the spiral ganglion

Species	No. of specimens	No. of cells	Turns of the cochlea
<i>Rhinolophus ferrumequinum</i>	2	15,724	3½
<i>Rhinolophus hipposideros</i>	5	13,674	3½
<i>Plecotus austriacus</i>	1	15,236	2½
<i>Cavia porcellus</i>	5	19,323	4½
<i>Felis silvestris</i>		38,760	3
<i>Homo sapiens</i>		29,024	2½
<i>Talpa europaea</i>	3	5,444	2½

The numbers of *Cavia*, *Felis* and man are taken from Firbas, Wicke & Volavsek (in press), Schuknecht (1960), and Guild (1932). After comparing these numbers, it can be inferred, that the number of nerve cells in the spiral ganglion depends on the dimension of the cochlea, and on

the range of perceived sounds. The neurons in the basal coil are responsible for the perception of high frequency sounds. It follows that in both the mole and the bat, the number of cells in this coil corresponds to the frequency range. The Talpidae are placed by Henson in the low frequency group. Our own investigation of the mole's cochlea also indicates the special sensitivity of this animal to low sounds. A study of the distribution of the nerve cells to the different coils revealed interesting details. It was seen, that in the Rhinolophidae the percentage of cells belonging to the basal turn is a little lower than in *Plecotus*, but nevertheless much higher than in the guinea pig. In *Rhinolophus* 38 % of the total number of cells belongs to the basal coil, in *Plecotus* 48 % and in *Cavia* 22 %. These percentages depend on the size of the basal coil, on the arrangement of cells, and on the number of turns. Nevertheless the highest absolute values are found in bats for cochlea of that size. As usual the spiral ganglion does not extend to the apex of the cochlea. Therefore in *Rhinolophus* the ganglion has three turns instead of three and a half in Corti's organ.

To get a better idea of the arrangement of cells, the density of cells in the different coils was determined. On photographs the nerve cells were counted on sections through the ganglion of a known area. This revealed an interesting difference between the Rhinolophidae and Vespertilionidae. The density increases in Rhinolophidae from basal to apical. In Vespertilionidae and in one species of Phyllostomatidae (*Phyllostomus hastatus*) the density is nearly equal in the basal and the upper coil. In table II the densities per 0.01 mm² of the ganglion in *Rhinolophus* and in *Cavia* are compared. The high density in *Rhinolophus* is obvious. The diameter of spiral ganglion cell nuclei is smaller in bats: 6 μ against 8.3 in *Cavia*.

Table II. *Density of ganglion cells in spiral ganglion.*

Species	Ganglion cells/0.01 mm ² of section through spiral ganglion		
	Basal	Middle	Apical
<i>Rhinolophus ferrumequinum</i>	62	75	87
<i>Rhinolophus hipposideros</i>	62	76	83
<i>Cavia porcellus</i>	32	33	30
			31

Following the procedure of Balogh, the acetylcholinesterase activity was used to show up the efferent bundle in the cochlear nerve. The method of Karnovsky & Roots was applied to demonstrate the acetylcholinesterase in the efferent fibres of the Rasmussen's bundle. In *Rhinolophus hipposideros* a lot of efferent fibres were revealed by this technique. In the sections obtained, the efferent fibres were more diffusely arranged and seemingly more numerous than in the rat or the cat. But at present it is not certain whether there is a really large number of efferent fibres in the bats cochlea. In the guinea pig the number of efferent fibres is about 500. In the cat the distribution of these fibres in the cochlea is known (Ishii & Blogh, 1968). A similar investigation of the bat's cochlea is planned. Kimura could demonstrate efferent nerve endings on the hair cells of a bat.

A preliminary examination of hair cells in the organ of Corti was made. From two cochleae of *Rhinolophus ferrumequinum* surface preparations of Corti's organ were made with help of a microdissection. The cell nuclei were stained with gallo-cyanin-chromealum. An eye-piece micrometer was used to count the outer and inner hair cells per

Table III

Number of hair cells per mm length of the organ of Corti in different mammals

Species	No. of outer hair cells	No. of inner hair cells
<i>Rhinolophus ferrumequinum</i>	454 basal 453 middle 370 apical	196 basal 163 middle 150 apical
<i>Cavia porcellus</i>	346 basal 365 2nd turn 459 3rd turn 257 apical	106 basal 101 2nd turn 133 3rd turn 70 apical
<i>Felis silvestris</i> (cf. Schuknecht, 1960)	415	120
Man (cf. Bredberg, 1968)	290 basal 525 apical	80 basal 115 apical

mm length of the organ of Corti. Because of the small number of countings a statistical analysis could not be made. The results obtained are listed in table III. Some comparisons from the available literature are also given. The number of outer and especially of inner hair cells per mm length of Corti's organ is largest in the basal turn. This corresponds with the data of Henson (1969), who finds dense nerve fibres in the basal coil of *Rhinolophidae*, *Hipposideridae*, and of *Chilonycteris*. It is probable, that the ratio of hair cells to spiral ganglion cells in the basal coil is higher for bats than for other species. That would give other possibilities in coding the acoustic signals. There is still controversy about the relation of inner and outer hair cells to spiral ganglion cells. According to Spoendlin (1969) most of the ganglion cells in the cat belong to the inner hair cells. The ratio total number of hair cells to total number of ganglion cells is approximately 1 : 2 in guinea pig and man, and 1 : 3 in cat. The ratio for bats is not yet known.

Finally the numbers of nerve cells in the spiral ganglion and in the cochlear nuclei are compared (table IV).

Table IV

Number of neurons in the spiral ganglion and in the cochlear nuclei

Species	No. of neurons in the spiral ganglion	No. of neurons in the cochlear nuclei
<i>Plecotus</i>	15,236	40,000 (Hall, 1969)
<i>Macacus</i>	31,247	88,000 (Chow, 1951)
	(Gacek & Rasmussen, 1961)	

The ratio of approximately 1 : 3 is the same in ultrasound sensitive and echolocating bat as in the monkey. According to Hall (1967) this ratio in whales is 1 : 5 to 1 : 8. As it is emphasized by Lütgemeier (1962) and Zvorykin (1959) the sub-cortical parts of the auditory pathway (like olivary superior and colliculus inferior) are extraordinary well developed in bats. Probably the neurophysiological correlation of the ability of echolocation can be seen in these central parts of the auditory system. On the other hand the ear apparatus and the peripheral parts of the auditory pathway are adapted for the perception of ultrasound.

LITERATURE CITED

- BALOGH, K. JR. & Y. NOMURA, 1964. A technique for the demonstration of acetylcholinesterase activity in the inner ear after decalcification with EDTA. *J. histochem. Cytochem.*, **12**: 931—933.
- BREDBERG, G., 1968. Cellular pattern and nerve supply of the human organ of Corti. *Acta oto-laryng.*, Suppl.: 236.
- CHOW, K. L., 1951. Numerical estimates of the auditory central nervous system of the Rhesus monkey. *J. comp. Neurol.*, **95**: 159—175.
- FLÖDERUS, S., 1944. Untersuchungen über den Bau der menschlichen Hypophyse mit besonderer Berücksichtigung der quantitativen mikromorphologischen Verhältnisse. *Acta path. microbiol. scand.*, Suppl.: 53.
- FIRBAS, W., W. WICKE & CH. VOLAVSEK, in press. Über Zahl und Anordnung der Ganglienzellen im Ganglion spirale des Meerschweinchens. *Msch. Ohr. Laryng. Rhinol.*
- GACEK, R. R. & G. L. RASMUSSEN, 1961. Fiber analysis of stato-acoustic nerve of guinea pig, cat and monkey. *Anat. Rec.*, **139**: 455—463.
- GUILD, S. R., 1932. Correlation of histologic observations and the acuity of hearing. *Acta oto-laryng.*, **17**: 207—245.
- HALL, J. G., 1967. Hearing and primary auditory centres of the whales. *Acta oto-laryng.*, Suppl.: 224.
- , 1969. The cochlea and the cochlear nuclei in the bat. *Acta oto-laryng.*, **67**: 350—353.
- HENSON, O. W., JR., 1961. Some morphological and functional aspects of certain structures of the middle ear in bats and insectivores. *Univ. Kansas sci. Bull.*, **12**: 151—255.
- , 1969. Neuroanatomical and physiological correlates of acoustic orientation. *Anat. Rec.*, **163**: 305.
- ISHII, D. & K. BALOGH, JR., 1968. Distribution of efferent nerve endings in the organ of Corti. *Acta oto-laryng.*, **66**: 282—288.
- KARNOVSKY, H. J. & L. ROOTS, 1964. A 'direct-colouring' thiocholin method for cholinesterases. *J. histochem. Cytochem.*, **12**: 219—221.
- KIMURA, R. S., 1968. After discussion, in: J. D. PYE: Hearing in bats. In: *Hearing mechanisms in vertebrates*. Ciba Foundation Symp. (Churchill, London) 1962.
- LÜTGEMEIER, F., 1962. Histologische Sonderheiten der Gehirne von Chiropteren verschiedener Körpergröße und verschiedener Orientierungsweise. *Z. Morph. Ökol.*, **50**: 687—725.
- SCHUKNECHT, H. F., 1960. Neuroanatomical correlates of auditory sensitivity and pitch discrimination in the cat. In: *Neural mechanisms of the auditory and vestibular systems*. (Thomas, Springfield).
- SPOENDLIN, H., 1969. Innervation patterns in the organ of Corti of the cat. *Acta oto-laryng.*, **67**: 239—254.
- WÜSTENFELD, E., 1957. Experimentelle Untersuchungen zum Problem der Schallanalyse im Innenohr. *Z. mikros. anat. Forsch.*, **63**: 327—387.
- ZVORYKIN, V. P., 1959. Morphological basis of locative and supersonic abilities in bat. *Arch. Anat. Hist. Embr.*, **36** (5): 19—31.