Short notes and reviews

A *Storthyngura* (Crustacea, Isopoda) from the Bellingshausen Sea, Antarctica

Angelika Brandt

Zoological Institute and Zoological Museum, Martin-Luther-King-Platz 3, D-20146 Hamburg, Germany.

E-mail: Abrandt@zoologie.uni-hamburg.de

Keywords: Southern Ocean, Bellingshausen Sea, Isopoda, *Storthyngura*

Abstract

During the expedition ANT XI/3 with RV *Polarstern* 17 multiple corers were taken in the Bellingshausen Sea. From this material only a single, immature isopod was sorted. This specimen is a manca II of a presumably new species of the genus *Storthyngura*. However, as it is immature and only a single specimen was sampled, more material of this species will have to be collected prior to the description of a new species.

Introduction

In 1994 an expedition with RV *Polarstern* aimed to investigate an area of the Antarctic and Southern Ocean which was not very well known biologically and geologically: the Bellingshausen and the Amundsen Sea. During this expedition a variety of different gear on board RV *Polarstern* was used and analysed (Miller & Grobe, 1996). Seventeen cores of the multiple corer (MUC) and a surface area of 10x10cm² from giant-box corer samples were sorted for peracarid crustaceans but only a single isopod was found. It is a specimen belonging to the genus *Storthyngura* (Vanhöffen, 1914) which occurs primarily in the deep sea, and was also reported from the Southern Ocean with several species. The species, which seems to be most closely related to the specimen sampled with the MUC is *Storthyngura fragilis* (Beddard, 1885), which was found off Marion Island, but has a distribution extending to Yokohama, Japan.

Material and methods

During the expedition ANT XI/3 with RV *Polarstern* in 1994, the specimen illustrated in the present paper was collected by Dr. Franz Gingele by means of a MUC at station 2540-1 at 70°03.8'S 87°55.9'W, in 1822 m depth (17.2.1994) (Miller & Grobe, 1996). The surface sediment consisted of foraminiferal mud (24% CaCO₃, 28 % sand, 42 % silt; an 30 % clay). Moreover, the station was characterized by strong current velocities close to the bottom (Miller & Grobe, 1996).

On board the vessel the animals were fixed in formalin (4%) and later transferred into ethanol (70%). The drawings were prepared with the help of a camera lucida. The specimen is deposited in the Zoological Museum of Hamburg under the catalogue number ZMH 39506.

Results and discussion

From the expedition ANT XI/3 only a single isopod specimen was sampled by means of the MUC in 17 hauls, indicating that Isopoda are either quite rare in the Bellingshausen Sea or that they occur very patchily and were therefore only collected as a single individual in that area of the Southern Ocean. The specimen is a member of *Storthyngura*, a typical deep-sea genus of the Munnopsidae (Eurycopinae), which usually occur deeper than 1500 m depth, often deeper than 4000. *Storthyngura* is distributed world wide and is reported with 10 species from the Southern Ocean until now. The presumed new species is illustrated in dorsal and lateral view (Fig. 2). Until now, many other species have been reported from the Southern Ocean by Pfeffer, 1887, Brandt, 1991, and Winkler, 1994. However, of all species of *Storthyngura* only *S.*

Fig. 1. Map of the Bellingshausen Sea with location of the MUC station PS 2540; no depth data are available close to the coastline due to heavy sea ice.

robustissima Monod, 1925, has been sampled in the Bellingshausen Sea in 400–1455 m depth and *S. praegrandis* George & Menzies, 1968, is known from the Drake Passage in the vicinity of the Bellingshausen Sea. However, both of these species differ from the species illustrated here, in bearing two strong acute spines on the head, and a narrower and shorter pleotelson, with a shorter and broader acute distal apex.

The new species from the Bellingshausen Sea, resembles most closely *S. fragilis* (Beddard, 1885). This species was first sampled off Marion Island, Subantarctic, but has also been reported from Yokohama, Japan, in 3429 m depth (Wolff, 1962). Both, Beddard (1885) and Wolff (1962) have not presented a complete description of all appendages. For that reason the immature (manca II) *Storothyngura* specimen from the Bellingshausen Sea was compared with the paratypes of *S. fragilis* (Natural History Museum, London, 1958.10.14:4–5). The pleotelson of the Bellingshausen Sea

Storothyngura is much longer, the frontolateral, and caudolateral spines are longer and more acute, and also the caudal acuminate tip of the pleotelson is slightly more elongated. The first peduncular article of the antennula is distally more slender than in *S. fragilis*, it is more acuminate distally and extends to the tip of the second peduncular article; of the 10 flagellar articles the last one bears two long aesthetascs, a feather-like seta and a simple one. The left mandible bears a pars incisiva with 4 teeth, a lacinia mobilis with 3 teeth, and a spine row of 8, mostly setulated setae. The fifth pereopod of the Bellingshausen Sea specimen bears 6 long feather-like setae dorsally on the basis, whereas *S. fragilis* bears only a single seta.

Even if the differences concerning the pleotelson and the antennula between *S. fragilis* and the manca II from the Bellingshausen Sea would be due to allometry, it is quite unlikely that an immature would bear more feather-like setae on the basis of pereopod five than an adult. Therefore it is quite

Fig. 2. Manca II of *Storthyngura* sp. ZMH 39506

likely that the immature specimen illustrated here only in dorsal and lateral views is a new species. However, as only a single immature specimen is

available, we will have to await the collection of further specimens, before a complete description of this presumably new species will be possible.

Acknowledgements

The author is most grateful to Dr. Franz Gingele (IOW, Warnemünde), who kindly made the material of *Storothyngura* available, to Dr. Wolfgang Bonn, who fixed and processed the material on board of *Polarstern*, and to Ms Monika Hänel for inking the dorsal views of the illustrations from the pencil drawings. Dr. Claus-Dieter Hillenbrand supplied the station map. Moreover thanks are due to the crew of the RV *Polarstern* during the expedition ANT XI/3 as well as to the Alfred-Wegener-Institute for Polar- and Marine Research, for logistics. Ms. Miranda Lowe, British Museum of Natural History, London, kindly made the paratype of *Storothyngura fragilis* available.

References

- Brandt A. 1991. Zur Besiedlungsgeschichte des antarktischen Schelfes am Beispiel der Isopoda (Crustacea: Malacostraca). *Ber. Polarforsch.* 98: 1-240.
- Beddard FE. 1885. Preliminary Notice of the Isopoda collected during the Voyage of H. M. S Challenger. Part II Munnop-sidae. – *Proc. zool. Soc. Lond.* 1885: 916-925.
- George RY, Menzies RJ. 1968. Species of *Storothyngura* (Isopoda) from the Antarctic with description of six new species. *Crustaceana* 14: 275-301.
- Miller H, Grobe H. 1996. The expedition ANTARKTIS XI/3 of RV *Polarstern* in 1994. *Ber. Polarforsch.* 188.
- Monod T. 1925. Isopodes et Amphipodes de l'Expédition antarctique belge (s. y. Belgica). II. *Bull. Mus. Hist. Nat., Paris* 31: 296-299.
- Pfeffer G. 1887. Die Krebse von Süd-Georgien nach der Ausbeute der deutschen Station, 1882-83. *Jahrb. Hamb. Wiss. Anst.* 4: 44-150.
- Vanhöffen E. 1914. Die Isopoden der Deutschen Südpolar-Expedition 1901-1903. *Deutsche Südpolarexpedition 1901-1903*, 25 (Zoologie) 7: 447-598.
- Winkler H. 1994. Characterization of the Scotia Arc isopod fauna (Crustacea, Malacostraca) from a biogeographical point of view: A multivariate approach. *Ber. Polarforsch.* 139: 1-196.
- Wolff T. 1962. The systematics and biology of bathyal and abyssal Isopoda Asellota. *Galathea Rep.* 6: 1-320.

Received: 28 July 1999