REVIEW

Evolution and Diversification of Land Plants. K. Iwatsuki & P.H. Raven (eds.), xiii + 330 pp., illus., 1997. Springer-Verlag, Tokyo, Berlin, Heidelberg, New York (in cooperation with the Botanical Society of Japan), ISBN 4-431-70203-2. Price: ¥ 11,000 (hard cover). For members of the Botanical Society of Japan ¥ 6,000.

Although not dealing with wood anatomy (wood evolution and xylem structure are only mentioned in passing a few times) this collection of invited essays on the diversification of land plants will be of interest to many readers of the IAWA Journal. The chapters are arranged under three headings: (1) origin and diversification of primitive land plants, (2) origin and diversification of angiosperms, and (3) speciation and mechanisms of diversification. The 24 contributing authors are all well acknowledged authorities in their respective fields. Wood anatomists interested in the evolution of the early woody land plants, conifers and angiosperms may (as I did) find the following papers especially interesting: D. Edwards, 'Charting diversity in early land plants: some challenges for the next millennium', which is especially informative in summarising recent progress in our understanding; T.A. Ohsawa's detailed phylogenetic reconstruction of some conifer families based on seed cone characters (one is tempted to overlay his cladograms with wood anatomical features to improve our very poor understanding of wood anatomical evolution in conifers); P. Endress' chapter on evolutionary biology of flowers discussing key innovations in the angiosperm flower which now characterise major clades; E.M. Friis et al.'s 'Fossil history of Magnoliid angiosperms', giving a critical summary of what is known of the oldest fossils of each of the constituting orders; D.E. Soltis et al.'s survey of molecular phylogenetic relationships among angiosperms; and J.A. Wolfe's chapter on relations of environmental change to angiosperm evolution during the late Cretaceous and Tertiary, which should help to focus our minds on ecological adaptation and homoplasy in wood structure. This book thus offers a very welcome state of the art on modern research in plant evolution, combining paleobotany, phylogenetic analysis and speciation research.

[P. B.]

ERRATUM — Following here is the correct legend of Fig. 42 on page 226 in the article by Sidiyasa & Baas, IAWA Journal 19 (2): 207–229.

Fig. 42. Transformations of wood anatomical characters of the genus *Alstonia*.

1. Average vessel frequency: ● = 3–10(–15); ○ = 29–61; ■ = (72–)100–149.
2. Average vessel diameter: ● = (76–)92–162 µm; ○ = 49–91 µm; ■ = 41–59 µm.
3. Axial parenchyma: ● = densely reticulate and with short or discontinuous lines, mostly uniseriate; ○ = reticulate with mostly irregularly spaced bands, 1–3(–4) cells wide; ■ = predominantly diffuse and diffuse-in-aggregates.
4. Fibre walls: ● = thin or very thin; ○ = medium to thick.
5. Fibre pits: ● = common in both radial and tangential sections; ○ = very rare in tangential section.
6. Latex tubes or laticifers: ● = present; ○ = absent.