REVIEW

Anatomía de maderas de México: Especies de una selva alta perennifolia I.

Josefina Barajas-Morales, Guillermo Ángeles Álvarez & Patricia Solís Sánchez, 126 pp., illus., 1997. Publicaciones Especiales del Instituto de Biología No. 16. D.R. Universidad Nacional Autónoma de México, Instituto de Biología, Apartado postal 70-223, 04510 México, DF. ISBN 968-36-6547-0. Price: US\$ 50.

This publication continues a series describing woods from various forest regions of Mexico. Sixty species (representing 55 genera in 30 families) from the tropical rain forest region of Los Tuxtlas in Veracruz are described. Common names, distribution, bark characteristics, and wood uses are given in addition to macroscopic and microscopic wood descriptions. Each description is accompanied by a color photograph of a board (nice reproduction), and four black-and-white photographs (2 transverse, 1 radial, 1 tangential). General features (e.g., color, odor, taste, specific gravity) and anatomical features are summarized in two tables. Atlases, such as this one, provide useful data on the diversity of wood structure found in different forest types, and are valuable references helpful for wood identification and determining the characteristics of particular taxa. My Spanish is definitely not the best, but the introduction to the atlas indicates that while the IAWA Feature List was followed for qualitative features, for quantitative features it was not. Vessel frequency was determined by counting each vessel group as one, i.e., a radial multiple of four counted as one unit, rather than as four as would result from counting each vessel individually. For vessel tangential diameter only the larger vessels were measured ("considerando sólo los vasos de diámetro mas grande"). Data on vessel diameter and frequency from this publication will not be comparable with data gathered by counting each vessel individually and measuring all width classes of vessels. The magnifications chosen for the tangential and radial sections of all species are fairly high, so while presence/absence of septate fibres and crystals can be seen, for some species a lower magnification tangential view would have permitted a clearer view of ray structure. Generally, the contrast of the photographs allows appreciating structural details, but there are some photographs that would have benefited from being printed darker and at a higher contrast, e.g., radial view of *Meliosma dentata* (Liebm.) Urban where the bars of the perforations are very, very faint. Nonetheless, the publication represents considerable work and will be a useful reference for years to come.

Elisabeth Wheeler