

Relationship between new farming practices and chemical use and the consumption of giant cricket (*Brachytrupes membranaceus* Drury, 1770)

F.O. Miantzia, F. Meutchieye and S. Niassy

Table S1. Respondents' age group age and gender structure.

Age group	Female	Male	Subtotal
10-20	3	16	19
20-30	15	64	79
30-40	24	57	81
40-50	30	56	86
50-60	11	33	44
60-70	3	7	10
Subtotal	86	233	319

Table S2. List of common weeds encountered in farmers' plots and use of herbicides in the Cameroon west region.

Family (number of species)	Scientific names
Amaranthaceae (2)	<i>Amaranthus viridis</i> Linnaeus <i>Amaranthus spinosus</i> Linnaeus
Asteraceae (5)	<i>Acanthospermum hispidum</i> dc <i>Ageratum conyzoides</i> Linnaeus Subsp. <i>Amaranthus spinosus</i> Linnaeus <i>Aspilia bussei</i> O. Hoffrn. et Muschl. <i>Bidens pilosa</i> Linnaeus
Commelinaceae (1)	<i>Commelina benghalensis</i> Linnaeus
Cyperaceae (3)	<i>Cyperus rotundus</i> Linnaeus <i>Cyperus esculentus</i> Linnaeus <i>Cyperus difformis</i> L.
Euphorbiaceae (2)	<i>Euphorbia heterophylla</i> Linnaeus <i>Euphorbia hirta</i> Linnaeus
Poaceae (16)	<i>Axonopus compressus</i> (sw.) P.Beauv. <i>Brachiaria ramosa</i> (Linnaeus) Stapf <i>Cenchrus biflorus</i> Roxb. <i>Chloris pilosa</i> Schumach. <i>Cynodon dactylon</i> (Linnaeus) Pers. <i>Digitaria horizontalis</i> Willd. <i>Echinochloa colona</i> (Linnaeus) Link <i>Eleusine indica</i> (Linnaeus) Gaertn. <i>Eragrostis aspera</i> (Jacq.) Nees <i>Eragrostis tremula</i> hochst. Ex steud. <i>Imperata cylindrica</i> (Linnaeus) P.Beauv. <i>Panicum laxum</i> sw. <i>Paspalum conjugatum</i> Berg. <i>Paspalum scrobiculatum</i> Linnaeus <i>Pennisetum polystachyon</i> (Linnaeus) Schult. <i>Setaria barbata</i> (Lam.) Kunth
Rubiaceae (1)	<i>Spermacoce chactocyhala</i> de Candolle
Tiliaceae (1)	<i>Triumpheta pentandra</i> A. Richard
Verbenaceae (1)	<i>Stachytarpheta angustifolia</i>